

Overfertilization, salt water, or road salt are often sources of salt.

CORRECTING DEFICIENCIES OR EXCESSES

Based on the soil test, applications of limestone, fertilizer, compost, or manure are often suggested. The proper time for application is also stated. Organic amendments are suggested when requested.

Liming materials are used to raise the pH of the soil. Dolomitic limestone is the most common form of liming material sold in Connecticut. It contains both calcium and magnesium carbonates. Quality limestone is finely ground to permit rapid release in the soil. Finely-ground limestone may be pelletized for use in rotary spreaders.

Hydrated lime reacts more rapidly but is somewhat caustic and may damage plants. If it is used, it should be thoroughly worked into the soil at about three-fourths the rate suggested for limestone. Burned lime or quicklime is very caustic and is rarely used.

The principal plant nutrients in mixed fertilizers are nitrogen (N), phosphorus (P), and potassium (K). Although they may be present in various forms, Connecticut law requires that the formula on the container be expressed as the percent of nitrogen (N), phosphoric acid (P_2O_5), and potash (K_2O). It is always in this order: thus a 5-10-5 fertilizer contains 5 pounds of N, 10 pounds of P_2O_5 , and 5 pounds of K_2O in each 100 pounds of fertilizer.

Trace elements are sometimes added to fertilizers. The desirable range between element toxicity and deficiency is usually narrow, so such mixtures may harm plants if used in excess.

If manure is applied, less commercial fertilizer may be required. Cow manure is low in nutrients and has a typical analysis of 0.5-0.25-0.5. About 10 tons per acre (1 cubic yard per 1000 square feet) may be applied. Fresh chicken manure

contains more nitrogen than cow manure, particularly in the ammonia form. If the plants to be grown are sensitive to ammonia, fresh manure should be aged, composted or worked into the soil well in advance of planting.

Other nitrogenous wastes such as municipal and industrial composts may be used. Application rates for composts are generally based on their nitrogen content, which is similar to animal manures. Use of composted sewage sludge may require approval of the Connecticut Department of Environmental Protection.

SOIL TESTING TELEPHONE NUMBERS

New Haven	(203) 974-8521
Windsor	(860) 683-4977

THE CONNECTICUT AGRICULTURAL EXPERIMENT STATION (CAES) prohibits discrimination on the basis of race, color, ancestry, national origin, sex, religious creed, age, political beliefs, sexual orientation, criminal conviction record, genetic information, learning disability, present or past history of mental disorder, mental retardation or physical disability including but not limited to blindness, or marital or family status. To file a complaint of discrimination, write Director, The Connecticut Agricultural Experiment Station, P.O. Box 1106, New Haven, CT 06504, or call (203) 974-8440. CAES is an equal opportunity provider and employer. Persons with disabilities who require alternate means of communication of program information should contact the Chief of Services at (203) 974-8442 (voice); (203) 974-8502 (FAX); or Michael.Last@po.state.ct.us.

SOIL TESTING



History of the Morgan Soil Test

Sampling the Soil

Tests Performed

Correcting Deficiencies or Excesses

**The Connecticut
Agricultural Experiment Station**

New Haven and Windsor

www.ct.gov/caes

HISTORY OF THE MORGAN SOIL TEST

More than 70 years ago, Dr. M.F. Morgan of this Station added immeasurably to our knowledge of the relationships between plants and soils. He devised a soil test that could estimate deficiencies or excesses of plant nutrients. Called the "Morgan Soil Test," it became the world's first widely accepted method for quickly estimating soil fertility.

Today, soil testing extends our knowledge of Connecticut soils and helps farmers, gardeners, and homeowners learn how to improve soil fertility in an environmentally responsible manner.

TESTS PERFORMED

Soil samples are tested for texture, organic matter, pH, nitrate nitrogen, ammonium nitrogen, phosphorus, potassium, calcium, and magnesium. Except for pH and texture, all results are expressed as high, medium, and low. A nutrient is classified as excessive when it is likely to damage plants. If necessary, we can perform tests for salts, micronutrients, and contaminants.

Texture: Texture of the soil influences the amount of water and nutrients a soil can hold. Sands, loamy sands, and sandy loams require more frequent watering and fertilizing than do fine sandy loams, loams and silt loams. Silty clay loams and clay loams drain poorly.

pH: Soil pH affects the availability of plant nutrients. When interpreted with texture and organic matter, pH indicates the lime needs of the soil. Acidic soils have a pH below 7.0 and alkaline soils have a pH above 7.0. Most plants grow best at a soil pH between 6.0 and 7.0. A small number of plants such as azalea, rhododendron, and blueberry prefer a soil pH between 4.5 and 5.5.

Nitrate and Ammonium Nitrogen tests indicate nitrogen immediately available to plants, but do not necessarily indicate how much nitrogen may later be liberated from the soil. Proper interpretation of these results is essential. Nitrogen favors leaf growth and imparts a deep green color to plant foliage. Excessive ammonium nitrogen can

damage plants and is often an indication of overfertilization. Very high nitrate nitrogen levels may increase the risk of nitrate contamination of surface and groundwater.

Phosphorus binds strongly in soil and is often unavailable to plants. Deficiencies in phosphorus are often indicated by poor root, fruit or vegetable growth and purpling of the older leaves. Excessive phosphorus can move to rivers, ponds and lakes and promote the growth of algae and weeds.

Potassium is supplied by the clay and organic matter in native soil. Improved plant growth is often obtained by the addition of potassium fertilizer. Sufficient potassium is thought to improve flowering, disease resistance, cold hardiness, and drought survival. Potassium leaches readily from soil.

Calcium in soil is readily revealed by our test.

Limestone can correct a calcium deficiency, and will also neutralize soil acidity. When the pH is correct and the calcium level is low, addition of gypsum (calcium sulfate) is often suggested.

Magnesium tests identify soils where dolomitic limestone or epsom salts (magnesium sulfate) are likely to be beneficial. A low magnesium level is usually associated with acidic soil.

Other Elements. Plants require small amounts of other elements including iron, copper, zinc, sulfur, and boron. Soil tests have been devised for these elements but they are neither quick nor infallible. Iron, copper, and zinc are affected by soil acidity and their availability can usually be inferred from the pH test. Sulfur is rarely deficient. Boron deficiency is usually encountered only where soils have a pH over 7.0. Measurements of soluble salts are sometimes reported on the tests.

SAMPLING THE SOIL

1. With a trowel, shovel, or auger, take thin slices or borings of soil from many places in the area to be sampled. Sample to a depth of 5 to 6 inches in gardens; 3 to 5 inches in lawns or permanent pastures.
2. Mix the soil thoroughly and place one pint of the mixture in a sealed plastic bag. Print your name, address and what you want to grow on a label and attach it to the outside of the bag. If any plant problems have occurred briefly describe them. State if you would like organic plant care suggestions.
3. If soils, fertilizer treatments, lime additions have been different, sample each area separately. Do not combine soil samples from different areas.
4. If the sample is very wet, let it dry before you submit it.
5. For more information and soil submission form (not required) on the internet go to <http://www.ct.gov/caes/soiltesting>.

Deliver or mail soil samples to Soil Tests, The Connecticut Agricultural Experiment Station, 123 Huntington Street, Box 1106, New Haven, CT 06504-1106 or 153 Cook Hill Road, Box 248, Windsor, CT 06095-0248.